

propofol injection. In addition, the tracking of alpha band SC values through operation will surely increase the possibility of patient awareness detection and prevent during and post-operative disturbance.

5 References

- [1] J.Bruhn, Depth of anaesthesia monitoring: what's available, what's validated and what's next?, *British Journal of Anaesthesia* 97: 85-94, 2006.
- [2] Jameson, L., Using EEG to monitor anaesthesia drug effects during surgery, *Journal of Clinical Monitoring and Computing* 20:445-472, 2006.
- [3] Ye, S., Kim J., Development for the evaluation Index of an Anaesthesia Depth Using the Bispectrum Analysis, *International Journal of Biological and Medical Sciences*, Vol. 1, No. 2, 2008.
- [4] Stockwell, R.G., Localization of the Complex Spectrum: The S Transform. *IEEE Transactions on Signal Processing*, Vol:44,No:4, 998-1001, April 1996.
- [5] Liu, C., Stockwell Coherence and its Application in Studying Motor Cortices under the Multi-Source Interference Task, *IEEE*, 2009.
- [6] Gürkan, G., Uslu, A., Akan, A., Demiralp, T., Topographic and Temporal Spectral Analysis of EEG Signals During Anaesthesia, 15th National Biomedical Conference, Antalya, Turkey, 21-24 April 2010.