

References

- [1] Jackson A, Buckley D, Parker GJM, Service SO. Dynamic contrast-enhanced magnetic resonance imaging in oncology. Springer; 2005. ISBN: 978-3-540-42322-5
- [2] Garpebring A, Ostlund N, Karlsson M. A Novel Estimation Method for Physiological Parameters in Dynamic Contrast-Enhanced MRI: Application of a Distributed Parameter Model Using Fourier-Domain Calculations [Article]. IEEE TRANSACTIONS ON MEDICAL IMAGING. 2009 SEP;28(9).
- [3] Koh TS, Zeman V, Darko J, Lee TY, Milosevic MF, Haider M, et al. The inclusion of capillary distribution in the adiabatic tissue homogeneity model of blood flow. Physics in medicine and biology. 2001 May;46(5):1519-1538.
- [4] Zhu XP, Li KL, Kamaly-Asl ID, Checkley DR, Tessier JLL, Waterton JC, et al. Quantification of endothelial permeability, leakage space, and blood volume in brain tumors using combined T1 and T2*contrast-enhanced dynamic MR imaging [Article]. JOURNAL OF MAGNETIC RESONANCE IMAGING. 2000 JUN;11(6).
- [5] Jiřík R, Bartoš M, Standara M, Taxt T: Blind Multichannel Estimation of Arterial Input Function in Dynamic Contrast-Enhanced MRI. Analysis of Biomedical Signals and Images, BIOSIGNAL 2010, in print.
- [6] Kershaw LE, Buckley DL. Precision in measurements of perfusion and microvascular permeability with T_1 -weighted dynamic contrast-enhanced MRI. Magnetic Resonance in Medicine. 2006;56(5):986-992.
- [7] Henderson E, Sykes J, Drost D, Weinmann HJ, Rutt BK, Lee TY. Simultaneous MRI measurement of blood flow, blood volume, and capillary permeability in mammary tumors using two different contrast agents. Journal of magnetic resonance imaging: JMRI. 2000 December;12(6):991-1003.